

# W15 Notes

## Complex algebra

---

Videos, Organic Chemistry Tutor

- [Complex numbers basics](#)

### 01 Theory - Complex arithmetic

The complex numbers  $\mathbb{C}$  are sums of real and imaginary numbers. Every complex number can be written uniquely in ‘Cartesian’ form:

$$z = a + bi, \quad a, b \in \mathbb{R}$$

To add, subtract, scale, and multiply complex numbers, treat ‘ $i$ ’ like a constant.

Simplify the result using  $i^2 = -1$ .

For example:

$$\begin{aligned} (1 + 3i)(2 - 2i) &\gg\gg 2 - 2i + 6i - 6i^2 \\ &\gg\gg 2 + 4i - 6(-1) \gg\gg 8 + 4i \end{aligned}$$


---

### ▣ Complex conjugate

Every complex number has a **complex conjugate**:

$$z = a + bi \gg\gg \bar{z} = a - bi$$

For example:

$$\overline{2 + 5i} = 2 - 5i$$

$$\overline{2 - 5i} = 2 + 5i$$

In general,  $\bar{\bar{z}} = z$ .

Conjugates are useful mainly because they eliminate imaginary parts:

$$(2 + 5i)(2 - 5i) \gg\gg 4 + 25 \gg\gg 29$$

In general:

$$(a + bi)(a - bi) \gg\gg a^2 - abi + bia - b^2i^2 \gg\gg a^2 + b^2 \in \mathbb{R}$$


---

### ⌚ Complex division

To divide complex numbers, use the conjugate to eliminate the imaginary part in the denominator.

For example, reciprocals:

$$\begin{aligned} \frac{1}{a+bi} &\gg\gg \frac{1}{a+bi} \cdot \frac{a-bi}{a-bi} \\ \gg\gg \frac{a-bi}{a^2+b^2} &\gg\gg \left(\frac{a}{a^2+b^2}\right) + \left(\frac{-b}{a^2+b^2}\right)i \end{aligned}$$

More general fractions:

$$\begin{aligned} \frac{a+bi}{c+di} &\gg\gg \frac{a+bi}{c+di} \cdot \frac{c-di}{c-di} \\ \gg\gg \frac{ac+bd+(bc-ad)i}{c^2+d^2} &\gg\gg \frac{ac+bd}{c^2+d^2} + \frac{bc-ad}{c^2+d^2}i \end{aligned}$$

### Multiplication preserves conjugation

For any  $z, w \in \mathbb{C}$ :

$$\overline{zw} = \bar{z}\bar{w}$$

Therefore, one can take products or conjugates in either order.

## 02 Illustration

### Example - Complex multiplication

Compute the products:

(a)  $(1-i)(4-7i)$     (b)  $(2+5i)(2-5i)$

#### Solution

(a)  $(1-i)(4-7i)$

Expand:

$$(1-i)(4-7i) \gg\gg 4-7i-4i+7i^2$$

Simplify  $i^2$ :

$$\gg\gg 4-7i-4i+7(-1)$$

$$\gg\gg -3-11i$$

(b)  $(2+5i)(2-5i)$

Expand:

$$(2+5i)(2-5i) \gg\gg 4-10i+10i-25i^2$$

Simplify  $i^2$ :

$$\gg\gg 4-10i+10i-25(-1) \gg\gg 29$$

### Example - Complex division

Compute the following divisions of complex numbers:

$$(a) \frac{1}{-3+i} \quad (b) \frac{1}{i} \quad (c) \frac{1}{7i} \quad (d) \frac{2+5i}{2-5i}$$

**Solution**

$$(a) \frac{1}{-3+i}$$

Conjugate is  $-3-i$ :

$$\frac{1}{-3+i} \gg \gg \frac{1}{-3+i} \cdot \frac{-3-i}{-3-i}$$

Simplify:

$$\gg \gg \frac{-3-i}{9+1} \gg \gg \frac{-3}{10} + \frac{-1}{10}i$$


---

$$(b) \frac{1}{i}$$

Conjugate is  $-i$ :

$$\frac{1}{i} \gg \gg \frac{1}{i} \cdot \frac{-i}{-i} \gg \gg -i$$


---

$$(c) \frac{1}{7i}$$

Factor out the  $1/7$ :

$$\frac{1}{7i} \gg \gg \frac{1}{7} \cdot \frac{1}{i}$$

Use  $\frac{1}{i} = -i$ :

$$\gg \gg \frac{1}{7} \cdot (-i) \gg \gg \frac{-1}{7}i$$


---

$$(d) \frac{2+5i}{2-5i}$$

Denominator conjugate is  $2+5i$ :

$$\frac{2+5i}{2-5i} \gg \gg \frac{2+5i}{2-5i} \cdot \frac{2+5i}{2+5i}$$

Simplify:

$$\gg \gg \frac{4+20i+25i^2}{4+25} \gg \gg \frac{-21}{29} + \frac{20}{29}i$$

## Complex exponential

Videos, Khan Academy

- [Complex exponential form](#)

### 03 Theory - cis, Euler, products, powers

Multiplication of complex numbers is much easier to understand when the numbers are written using polar form.

There is a shorthand ‘cis’ notation:

$$\begin{aligned} a + bi &\gg \gg r \cos \theta + r \sin \theta i \\ &\gg \gg r(\cos \theta + i \sin \theta) \\ &\gg \gg r \operatorname{cis} \theta \end{aligned}$$

The cis notation stands for  $\cos \theta + i \sin \theta$ .

For example:

$$\begin{aligned} \sqrt{2} - \sqrt{2}i &\gg \gg 2\left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i\right) \\ &\gg \gg 2 \cos\left(-\frac{\pi}{4}\right) + 2 \sin\left(-\frac{\pi}{4}\right)i \\ &\gg \gg 2 \operatorname{cis}\left(-\frac{\pi}{4}\right) \end{aligned}$$


---

### Euler Formula

General **Euler Formula**:

$$r e^{i\theta} = r \cos \theta + i r \sin \theta$$

On the unit circle:

$$e^{i\theta} = \cos \theta + i \sin \theta$$

The form  $r e^{i\theta}$  expresses the *same data* as the cis form.

The principal advantage of the form  $r e^{i\theta}$  is that it *reveals the rule for multiplication*:

### Complex multiplication - Exponential form

$$r_1 e^{i\theta_1} \cdot r_2 e^{i\theta_2} = (r_1 r_2) e^{i(\theta_1 + \theta_2)}$$

In words:

- Multiply radii
- Add angles

Notice:

$$\text{multiply by } e^{i\frac{\pi}{2}} \iff \text{rotate by } +90^\circ$$

Notice:

$$e^{i\frac{\pi}{2}} = +i$$

Therefore  $i$  ‘acts upon’ other numbers by rotating them  $90^\circ$  counterclockwise!

---

### De Moivre's Theorem - Complex powers

In exponential notation:

$$(re^{i\theta})^n = r^n e^{i \cdot n\theta}$$

In cis notation:

$$(r \operatorname{cis} \theta)^n = r^n \operatorname{cis}(n\theta)$$

Expanded cis notation:

$$(r \cos \theta + i r \sin \theta)^n = r^n \cos(n\theta) + i r^n \sin(n\theta)$$

So the power of  $n$  acts like this:

- **Stretch:**  $r$  to  $r^n$
- **Rotate** by  $n$  increments of  $\theta$

### ☰ Extra - Derivation of Euler Formula

Recall the power series for  $e^x$ :

$$e^x = 1 + \frac{1}{1!}x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \dots = \sum_{i=0}^{\infty} \frac{1}{i!}x^i$$

Plug in  $x = i\theta$ :

$$e^{i\theta} \ggg 1 + (i\theta) + \frac{1}{2!}(i\theta)^2 + \frac{1}{3!}(i\theta)^3 + \dots$$

Simplify terms:

$$\ggg 1 + i\theta - \frac{1}{2!}\theta^2 - \frac{1}{3!}i\theta^3 + \frac{1}{4!}\theta^4 + \frac{1}{5!}i\theta^5 - \frac{1}{6!}\theta^6 - \frac{1}{7!}i\theta^7 + \frac{1}{8!}\theta^8 + \dots$$

Separate by  $i$ -factor. Select out the  $\overbrace{\text{terms with } i}$ :

$$\ggg 1 + \overbrace{i\theta} - \frac{1}{2!}\theta^2 - \frac{1}{3!}\overbrace{i\theta^3} + \frac{1}{4!}\theta^4 + \frac{1}{5!}\overbrace{i\theta^5} - \frac{1}{6!}\theta^6 - \frac{1}{7!}\overbrace{i\theta^7} + \frac{1}{8!}\theta^8 + \dots$$

Separate into a series without  $i$  and a series with  $i$ :

$$\ggg \left(1 - \frac{1}{2!}\theta^2 + \frac{1}{4!}\theta^4 - \dots\right) + \left(\theta - \frac{1}{3!}\theta^3 + \frac{1}{5!}\theta^5 - \dots\right)i$$

Identify  $\cos \theta$  and  $\sin \theta$ . Write trig series:

$$\cos \theta = 1 - \frac{1}{2!}\theta^2 + \frac{1}{4!}\theta^4 - \dots$$

$$\sin \theta = \theta - \frac{1}{3!}\theta^3 + \frac{1}{5!}\theta^5 - \dots$$

Therefore  $e^{i\theta} = \cos \theta + i \sin \theta$ .

## 04 Illustration

### ☰ Example - Complex product, quotient, power using Euler

Start with two complex numbers:

$$z = 2e^{i\frac{\pi}{2}} \quad w = 5e^{i\frac{\pi}{3}}$$

Product  $zw$ :

$$\begin{aligned} zw &\gg (2e^{i\frac{\pi}{2}}) \cdot (5e^{i\frac{\pi}{3}}) \\ &\gg (2 \cdot 5) (e^{i\frac{\pi}{2}}) (e^{i\frac{\pi}{3}}) \gg 10e^{i\frac{\pi}{2} + i\frac{\pi}{3}} \gg 10e^{i\frac{5\pi}{6}} \end{aligned}$$

Quotient  $z/w$ :

$$\begin{aligned} z/w &\gg \frac{(2e^{i\frac{\pi}{2}})}{(5e^{i\frac{\pi}{3}})} \\ &\gg \frac{2e^{i\frac{\pi}{2}}}{5e^{i\frac{\pi}{3}}} \gg \frac{2}{5} e^{i\frac{\pi}{2}} e^{-i\frac{\pi}{3}} \gg \frac{2}{5} e^{i\frac{\pi}{6}} \end{aligned}$$

Power  $z^8$ :

$$\begin{aligned} z^8 &\gg (2e^{i\frac{\pi}{2}})^8 \\ &\gg 2^8 (e^{i\frac{\pi}{2}})^8 \gg 512e^{i \cdot 8\pi} \end{aligned}$$

Notice:

$$e^{i \cdot 4\pi} \gg (e^{2\pi i})^2 \gg 1^2 \gg 1$$

Simplify:

$$512e^{i \cdot 8\pi} \gg 512$$

Thus:  $z^8 = 512$ .

### Example - Complex power from Cartesian

Compute  $(3 + 3i)^4$ .

#### Solution

First convert to exponential form:

$$\begin{aligned} 3 + 3i &\gg 3\sqrt{2} \left( \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i \right) \\ &\gg 3\sqrt{2}e^{i\frac{\pi}{4}} \end{aligned}$$

Compute the power:

$$\begin{aligned} (3 + 3i)^4 &\gg (3\sqrt{2}e^{i\frac{\pi}{4}})^4 \\ &\gg 324e^{i\pi} \gg -324 \end{aligned}$$

## Complex roots

Videos, Trefor Bazett

- [Finding cube roots](#): Find cube roots of  $-1$

Videos, Brain Gainz

- [Finding nth roots](#): Fourth roots of  $\sqrt{3} - i$  and cube roots of  $-8$

## 05 Theory - Roots formula

The exponential notation leads to a formula for a complex  $n^{\text{th}}$  root of any complex number:

$$\sqrt[n]{re^{i\theta}} = \sqrt[n]{r} e^{i\frac{\theta}{n}}$$

⚠ Every complex number actually has  $n$  distinct complex  $n^{\text{th}}$  roots!

That's two square roots, three cube roots, four  $4^{\text{th}}$  roots, etc.

### ▣ All complex roots

The complex roots of  $z = re^{i\theta}$  are given by:

$$w_k = \sqrt[n]{r} \cdot e^{i\left(\frac{\theta}{n} + k\frac{2\pi}{n}\right)} \quad \text{for each } k = 0, 1, 2, \dots, n-1$$

In Cartesian notation:

$$w_k = \sqrt[n]{r} \cos\left(\frac{\theta}{n} + k\frac{2\pi}{n}\right) + \sqrt[n]{r} \sin\left(\frac{\theta}{n} + k\frac{2\pi}{n}\right)i$$

In words:

- Start with the basic root:  $\sqrt[n]{r} \cdot e^{i\frac{\theta}{n}}$
- Rotate by increments of  $\frac{2\pi}{n}$  to get all other roots

### ☰ Extra - Complex roots proof

We must verify that  $w_k^n = re^{i\theta}$ :

$$\left(\sqrt[n]{r} \cdot e^{i\left(\frac{\theta}{n} + k\frac{2\pi}{n}\right)}\right)^n \gg \gg r^{\frac{n}{n}} \cdot e^{i\left(\frac{\theta}{n} + k\frac{2\pi}{n}\right)n}$$

$$\gg \gg r \cdot e^{i(\theta + 2\pi k)} \gg \gg r e^{i\theta} e^{i2\pi k} \gg \gg r e^{i\theta}$$

## 06 Illustration

### ☰ Example - Finding all $4^{\text{th}}$ roots of 16

Compute all the  $4^{\text{th}}$  roots of 16.

#### Solution

Write  $16 = 16e^{0i}$ .

Evaluate roots formula:

$$(16e^{0i})^{\frac{1}{4}} \gg \gg w_k = 16^{\frac{1}{4}} e^{i\left(\frac{0}{4} + k\frac{2\pi}{4}\right)}$$

Simplify:

$$\gg \gg \quad 2e^{ik\frac{\pi}{2}} \quad \gg \gg \quad 2, 2i, -2, -2i$$

### ☰ Example - Finding 2<sup>nd</sup> roots of 2i

Find both 2<sup>nd</sup> roots of 2i.

#### Solution

Write  $2i = 2e^{i\frac{\pi}{2}}$ .

Evaluate roots formula:

$$(2e^{i\frac{\pi}{2}})^{\frac{1}{2}} \quad \gg \gg \quad w_k = \sqrt{2}e^{i\left(\frac{\pi/2}{2} + k\frac{2\pi}{2}\right)}$$

$$\gg \gg \quad \sqrt{2}e^{i\left(\frac{\pi}{4} + k\pi\right)}$$

Compute the options:  $k = 0, 1$ :

$$\gg \gg \quad \sqrt{2}e^{i\frac{\pi}{4}}, \sqrt{2}e^{i\frac{5\pi}{4}}$$

Convert to rectangular:


$$\gg \gg \quad \sqrt{2}\left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i\right), \sqrt{2}\left(-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i\right)$$

$$\gg \gg \quad 1+i, 1-i$$

### ☰ Example - Some roots of unity

Find the 1<sup>st</sup> and 2<sup>nd</sup> and 3<sup>rd</sup> and 4<sup>th</sup> and 5<sup>th</sup> and 6<sup>th</sup> roots of the number 1.

#### Solution



1<sup>st</sup>

Write  $1 = e^{0i}$ . Evaluate roots formula. There is no possible  $k$ :

$$(e^{0i})^{\frac{1}{1}} \quad \gg \gg \quad e^{0i} \quad \gg \gg \quad 1$$

2<sup>nd</sup>

Write  $1 = e^{0i}$ . Evaluate roots formula in terms of  $k$ :

$$(e^{0i})^{\frac{1}{2}} \gg \gg w_k = e^{i(\frac{0}{2} + k\frac{2\pi}{2})} \quad k = 0, 1$$

Compute the two options,  $k = 0, 1$ :

$$\gg \gg 1, e^{\pi i} \gg \gg 1, -1$$


---

3<sup>rd</sup>

Evaluate roots formula in terms of  $k$ :

$$(e^{0i})^{\frac{1}{3}} \gg \gg w_k = e^{i(\frac{0}{3} + k\frac{2\pi}{3})}$$

Compute the options:  $k = 0, 1, 2$ :

$$\gg \gg 1, e^{i\frac{2\pi}{3}}, e^{i\frac{4\pi}{3}} \gg \gg 1, -\frac{1}{2} + \frac{\sqrt{3}}{2}i, -\frac{1}{2} - \frac{\sqrt{3}}{2}i$$


---

4<sup>th</sup>

Evaluate roots formula:

$$(e^{0i})^{\frac{1}{4}} \gg \gg w_k = e^{i(\frac{0}{4} + k\frac{2\pi}{4})}$$

Compute the options:  $k = 0, 1, 2, 3$ :

$$1, e^{i\frac{\pi}{2}}, e^{i\pi}, e^{i\frac{3\pi}{2}} \gg \gg 1, i, -1, -i$$


---

5<sup>th</sup>

Evaluate roots formula:

$$(e^{0i})^{\frac{1}{5}} \gg \gg w_k = e^{i(\frac{0}{5} + k\frac{2\pi}{5})}$$

Compute the options:  $k = 0, 1, 2, 3, 4$ :

$$1, e^{i\frac{2\pi}{5}}, e^{i\frac{4\pi}{5}}, e^{i\frac{6\pi}{5}}, e^{i\frac{8\pi}{5}}$$

Don't simplify, it's not feasible.

---

6<sup>th</sup>

Evaluate roots formula:

$$(e^{0i})^{\frac{1}{6}} \gg \gg w_k = e^{i(\frac{0}{6} + k\frac{2\pi}{6})}$$

Compute the options:  $k = 0, 1, 2, 3, 4, 5$ :

$$1, e^{i\frac{2\pi}{6}}, e^{i\frac{4\pi}{6}}, e^{i\frac{6\pi}{6}}, e^{i\frac{8\pi}{6}}, e^{i\frac{10\pi}{6}}$$

Simplify:

$$\gg \gg 1, \frac{1}{2} + \frac{\sqrt{3}}{2}i, -\frac{1}{2} + \frac{\sqrt{3}}{2}i, -1, -\frac{1}{2} - \frac{\sqrt{3}}{2}i, \frac{1}{2} - \frac{\sqrt{3}}{2}i$$

