

W13 - Examples

Law of Large Numbers

Markov and Chebyshev

A tire shop has 500 customers per day on average.

- (a) Estimate the odds that more than 700 customers arrive today.
- (b) Assume the variance in daily customers is 10. Repeat (a) with this information.

Solution

Write X for the number of daily customers.

- (a) Using Markov's inequality with $c = 700$, we have:

$$P[X \geq 700] \leq \frac{500}{700} \approx 0.71$$

- (b) Using Chebyshev's inequality with $c = 200$, we have:

$$P[|X - 500| \geq 200] \leq \frac{100}{200^2} \approx 0.0025$$

The Chebyshev estimate is much smaller!

LLN: Average winnings

A roulette player bets as follows: he wins \$100 with probability 0.48 and loses \$100 with probability 0.52. The expected winnings after a single round is therefore $100 \cdot 0.48 - 100 \cdot 0.52$ which equals $-\$4$.

By the LLN, if the player plays repeatedly for a long time, he expects to lose \$4 per round on average.

The 'expects' in the last sentence means: the PMF of the cumulative average winnings approaches this PMF:

$$P_{M_n(X)}(k) = \begin{cases} 1 & k = \$4 \\ 0 & k \neq \$4 \end{cases}$$

This is by contrast to the 'expects' of expected value: the probability of achieving the expected value (or something near) may be low or zero! For example, a single round of this game.

Enough samples

Suppose $X_1, X_2, \dots$ are IID samples of $X \sim \text{Ber}(0.6)$.

- (a) Compute $E[X]$ and $\text{Var}[X]$ and $\text{Var}[M_{100}(X)]$.
- (b) Use the finite LLN to find α such that:

$$P[|M_{100}(X) - 0.6| \geq 0.05] \leq \alpha$$

(c) How many samples n are needed that to guarantee that:

$$P[|M_n(X) - 0.6| \geq 0.1] \leq 0.05$$

Statistical testing

One-tail test: Weighted die

Your friend gives you a single regular die, and say she is worried that it has been weighted to prefer the outcome of 2. She wants you to test it.

Design a significance test for the data of 20 rolls of the die to determine whether the die is weighted. Use significance level $\alpha = 0.05$.

Solution

Let X count the number of 2s that come up.

The Claim: “the die is weighted to prefer 2”

The null hypothesis H_0 : “the die is normal”

Assuming H_0 is true, then $X \sim \text{Bin}(20, 1/6)$, and therefore:

$$P_{X|H_0}(k) = \binom{20}{k} (1/6)^k (5/6)^{20-k}$$

⚠ Notice that “prefer 2” implies the claim is for *more* 2s than normal.

Therefore: Choose a one-tail rejection set.

Need r such that $P[X \geq r | H_0] = 0.05$

- Equivalently: $P[X < r | H_0] = 0.95$

Solve for r by computing conditional CDF values:

$k :$	0	1	2	3	4	5	6	7
$F_{X H_0}(k) :$	0.026	0.130	0.329	0.567	0.769	0.898	0.963	0.989

Therefore, choose $r = 6$. Then $P[X \geq r | H_0] < 0.04$ and no smaller (integer) r will produce significance below 0.05.

The final answer is:

$$R = \{x \mid x \geq 6\}$$

Two-tail test: Circuit voltage

A boosted AC circuit is supposed to maintain an average voltage of 130 V with a standard deviation of 2.1 V. Nothing else is known about the voltage distribution.

Design a two-tail test incorporating the data of 40 independent measurements to determine if the expected value of the voltage is truly 130 V. Use $\alpha = 0.02$.

Solution

Use $M_{40}(V)$ as the decision statistic, i.e. the sample mean of 40 measurements of V .

The Claim to test: μ is not 130

The null hypothesis H_0 : $\mu = 130$

Rejection region:

$$|M_{40} - 130| \geq c$$

where c is chosen so that $P[|M_{40} - 130| \geq c] = 0.02$

Assuming H_0 , we expect that:

$$E[M_{40}] = 130 \quad \sigma^2 = \text{Var}[M_{40}] = \frac{2.1^2}{40} \approx 0.110$$

Recall Chebyshev's inequality:

$$P[|M_{40} - 130| \geq c] \leq \frac{\sigma^2}{c^2} \approx \frac{0.110}{c^2}$$

Now solve:

$$\frac{0.110}{c^2} = 0.2 \quad \gg \gg \quad c \approx 0.74$$

Therefore the rejection region should be:

$$M_{40} < 129.26 \quad \cup \quad 130.74 < M_{40}$$

One-tail test with a Gaussian: Weight loss drug

Assume that in the background population in a specific demographic, the distribution of a person's weight W satisfies $W \sim \mathcal{N}(190, 24)$. Suppose that a pharmaceutical company has developed a weight-loss drug and plans to test it on a group of 64 individuals.

Design a test at the $\alpha = 0.01$ significance level to determine whether the drug is effective.

Solution

Since the drug is tested on 64 individuals, we use the sample mean $M_{64}(W)$ as the decision statistic.

The Claim: "the drug is effective in reducing weight"

The null hypothesis H_0 : "no effect: weights on the drug still follow $\mathcal{N}(190, 24)$ "

Assuming H_0 is true, then $W \sim \mathcal{N}(190, 24)$.

⚠ One-tail test because the drug is expected to *reduce* weight (unidirectional).

Rejection region:

$$M_{64}(W) \leq r$$

Compute $\frac{24}{\sqrt{64}} = 3$.

Since $W \sim \mathcal{N}(190, 24)$, we know that $M_{64}(W) \sim \mathcal{N}(190, 3^2)$.

Furthermore:

$$\frac{M_{64}(W) - 190}{3} \sim \mathcal{N}(0, 1)$$

Then:

$$\begin{aligned} P[M_{64}(W) < r] &= P\left[Z < \frac{r - 190}{3}\right] \\ &= \Phi\left(\frac{r - 190}{3}\right) \end{aligned}$$

Solve:

$$\begin{aligned} P[M_{64}(W) < r] &= 0.01 \\ \gg \gg \quad \Phi\left(\frac{r - 190}{3}\right) &= 0.01 \\ \gg \gg \quad \Phi\left(\frac{190 - r}{3}\right) &= 0.99 \\ \gg \gg \quad \frac{190 - r}{3} &= 2.33 \\ \gg \gg \quad r &= 183.01 \end{aligned}$$

Therefore, the rejection region:

$$M_{64}(W) \leq 183.01$$